

# **MICHTOX: A Mass Balance and Bioaccumulation Model for Toxic Chemicals in Lake Michigan**

**Ronald Rossmann, Editor**

Large Lakes and Rivers Forecasting Research Branch  
Mid-Continent Ecology Division  
National Health and Environmental Effects Research Laboratory  
Office of Research and Development  
Large Lakes Research Station  
9311 Groh Road  
Grosse Ile, Michigan 48138

**September 2005**



*Printed on Recycled Paper*

---

## **Notice**

The information in this document has been obtained primarily through funding by the United States Environmental Protection Agency (USEPA) under the auspices of the Great Lakes National Program Office (GLNPO) and the Office of Research and Development (ORD). The report has been subjected to the Agency's peer and administrative review and it has been approved for publication as a USEPA document. Mention of trade names or commercial products does not constitute endorsement or recommendation for use. Because the purpose of this report is to document the development of the MICHTOX model, conclusions are historical and should not be considered current.

---

## Foreword

Federal and contractor staff at the United States Environmental Protection Agency's Large Lakes Research Station have been involved with the development of mass balance models for the Great Lakes since the early 1970s. MICHTOX is a mass balance model developed to predict chemical concentrations in water and sediments of Lake Michigan in response to chemical loads to the lake. The model was adapted from the general water quality model WASP4. The MICHTOX bioaccumulation model was based upon the WASTOXv4 food chain model. Development of MICHTOX began in the early 1990s. The model was developed as a planning tool for the Lake Michigan Mass Balance Project (LMMBP) (U.S. Environmental Protection Agency, 1997). This work was documented in an in-house report in 1992 (Part 1). The model was applied as a screening-level model for atrazine in Lake Michigan in support of the LMMBP (Rygwelski *et al.*, 1999). The model was slightly revised and applied to polychlorinated biphenyls (PCBs) in Lake Michigan to confirm model results with the LMMBP project data and to provide preliminary modeling results for inclusion in the 2002 Lake-wide Management Plan (LaMP) report (Lake Michigan Technical Committee, 2002). These were reported in a 2002 contractor report (Part 2). The purpose of this report is to document through 2002 the progression of MICHTOX model development and application of the model to describing the behavior of contaminants, especially PCBs, in Lake Michigan. Both parts of this report have been cited numerous times in the literature. This report provides ready access to these for interested parties. For PCBs, results from application of the model have been superseded by more recent results.

Lake Michigan Technical Committee. 2002. Lake Michigan Lakewide Management Plan (LaMP), 2002. U.S. Environmental Protection Agency, Great Lakes National Program Office, Chicago, Illinois. 102 pp.

Rygwelski, K.R., W.L. Richardson, and D.D. Endicott. 1999. A Screening-Level Model Evaluation of Atrazine in the Lake Michigan Basin. *J. Great Lakes Res.*, 25(1):94-106.

U.S. Environmental Protection Agency. 1997. Lake Michigan Mass Budget/Mass Balance Work Plan. U.S. Environmental Protection Agency, Great Lakes National Program Office, Chicago, Illinois. EPA/905/R-97/018, 155 pp.

---

## **Abstract**

MICHTOX is a toxic chemical mass balance and bioaccumulation model for Lake Michigan. It was developed for the United States Environmental Protection Agency's Region V in support of the Lake Michigan Lake-wide Management Plan (LaMP) to provide guidance on expected water quality improvements in response to critical pollutant loading reductions. The 11 critical pollutants modeled were benzo(a)pyrene, chlordane, total dichlorodiphenyltrichloroethane (DDT), dieldrin, heptachlor epoxide, hexachlorobenzene, lead, total polychlorinated biphenyls (PCBs), 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), 2,3,7,8-tetrachlorodibenzofuran (TCDF), and toxaphene. Concentrations of these were predicted in 17 water and sediment segments in response to atmospheric and tributary loadings. The bioaccumulation model was coupled to the mass balance model to predict chemical accumulation in lake trout and bloater through pelagic and benthic food chains. Mass balance predictions were validated using plutonium, lead, and PCBs data; and bioaccumulation predictions were validated with PCBs data. The model was later applied to provide preliminary PCBs model results for the Lake Michigan Mass Balance Project. Results from this application were used to guide the development of a more resolute model for PCBs. Results for PCBs described in Part 1 are superseded by results in Part 2. Part 2 results have been replaced by a more recent application of MICHTOX that has been presented at various meetings and will be published at a future date. This document is meant to provide a historical perspective of MICHTOX development and application.

## Table of Contents

Notice .....	ii
Foreword .....	iii
Abstract .....	iv
Table of Contents .....	v
List of Figures .....	viii
List of Tables .....	xiv
Abbreviations .....	xvi
Acknowledgments .....	xvii
Executive Summary .....	xviii
 Part 1 1992 MICHTOX: A Mass Balance and Bioaccumulation Model for Toxic Chemicals in Lake Michigan .....	 1
1.1 Executive Summary .....	1
1.2 Recommendations .....	3
1.2.1 Verification of Model Predictions .....	3
1.2.1.1 Air Concentrations/Deposition Fluxes .....	3
1.2.1.2 Surficial Sediment Concentrations .....	3
1.2.1.3 Lake Trout .....	3
1.2.1.4 Major Tributaries .....	3
1.2.1.5 Water .....	3
1.2.2 Extend Model to Other Critical Pollutants and Target Organisms .....	3
1.2.3 Further Model Development .....	4
1.2.3.1 Circulation .....	4
1.2.3.2 Sediment Transport .....	4
1.2.3.3 Organic Carbon Dynamics .....	5
1.2.3.4 Food Chain Variability and Dynamics .....	5
1.2.4 Establish Linkages to Atmospheric and Watershed Models .....	5
1.3 Introduction .....	6
1.3.1 Project Objectives .....	6
1.3.2 Lake Michigan Toxics Problem .....	7
1.4 Model Description .....	8
1.4.1 Model Framework .....	8
1.4.2 Segmentation .....	10
1.4.3 Circulation .....	12
1.4.4 Solids Balance .....	13

1.4.5	Chemical Partitioning and Loss	15
1.4.5.1	Partitioning	15
1.4.5.2	Volatilization	16
1.4.5.3	Photolysis	18
1.4.5.4	Sediment-Water Diffusive Exchange	18
1.4.6	Chemical Loads and Boundary Conditions	18
1.4.6.1	Atmospheric Deposition	18
1.4.6.2	Tributary Loads	19
1.4.6.3	Loading Histories	20
1.4.6.3.1	Plutonium	20
1.4.6.3.2	Lead	20
1.4.6.3.3	PCBs	20
1.4.6.4	Lake Huron Boundary Conditions	21
1.4.7	Chemical Bioaccumulation	21
1.4.7.1	Food Chain	23
1.4.7.2	Uptake Rate	23
1.4.7.3	Elimination Rate	23
1.4.7.4	Dietary Accumulation	25
1.4.7.5	Modeling the Base of the Food Chain	25
1.4.8	Steady-State Model	26
1.4.9	Chemical-Specific Parameterization	28
1.5	Model Validation	30
1.5.1	Plutonium	31
1.5.2	Lead	32
1.5.3	PCBs	33
1.5.3.1	Water	33
1.5.3.2	Sediment	34
1.5.3.3	Biota	35
1.5.3.4	Bioaccumulation	38
1.6	Steady-State Model Applications	38
1.6.1	Steady-State Load-Response Predictions	41
1.6.2	Mass Fate and Transport	41
1.6.3	Sensitivity Analysis	44
1.6.4	Uncertainty Analysis	50
1.6.4.1	Analysis of Model Uncertainty	50
1.6.4.2	Results	51
1.6.4.3	Critical Parameterization Uncertainty	54
1.7	Dynamic Model Applications	57
1.7.1	Toxic Chemical Lag Time	57
1.7.2	PCBs Fate and Transport Fluxes in Dynamic Simulations	60
1.7.3	Additional Dynamic Simulations for PCBs	60
1.7.3.1	PCBs Control Scenarios	61
1.7.3.2	Severe Storm Event	61
1.7.4	Uncertainty in Dynamic Simulations	62
1.8	References	64
1.9	Appendix: Steady-State Model Output for Each Toxic Chemical	72

---

Part 2	2002 Lake Michigan Mass Balance Project: Modeling Total Polychlorinated Biphenyls Using the MICHTOX Model . . . . .	84
2.1	Executive Summary . . . . .	84
2.2	Recommendations . . . . .	85
2.3	Introduction . . . . .	85
2.4	Description of Model, Data, and Simulations . . . . .	86
2.4.1	MICHTOX Model . . . . .	86
2.4.2	Mass Balance Data for PCBs . . . . .	90
2.4.3	Revised MICHTOX and LMMBP Forcing Functions . . . . .	94
2.5	Results and Discussion . . . . .	98
2.5.1	Confirmation of MICHTOX PCBs Bioaccumulation Predictions . . . . .	98
2.5.2	Comparison of Original MICHTOX PCBs Simulations to the LMMBP Data . . . . .	104
2.5.3	Updating Parameterization for Henry's Constants . . . . .	109
2.5.4	Updating the Chemical Volatilization Rate Formulations . . . . .	110
2.5.5	Long-Term Hindcast/Forecast Simulations . . . . .	110
2.5.6	Toxic Chemical Management Forecasts . . . . .	129
2.5.7	Uncertainty of MICHTOX Model Predictions . . . . .	131
2.5.8	Are Lake Michigan Total PCBs Concentrations in Equilibrium With Atmospheric Vapor Concentrations? . . . . .	131
2.5.9	Sensitivity of Bioaccumulation Predictions to Initial Total PCBs Concentrations in Fish . . . . .	132
2.5.10	Sensitivity of Bioaccumulation Predictions to Food Chain Model Parameterization . . . . .	132
2.5.11	Are Total PCBs Bioaccumulation Factors Constant for Lake Trout in Lake Michigan? . . . . .	136
2.6	References . . . . .	136

---

## List of Figures

1.1	Long-term concentration trends for toxic chemicals in small Lake Michigan fish . . . . .	7
1.2	MICHTOX mass balance schematic . . . . .	9
1.3	Spatial segmentation for the 17 segment MICHTOX model . . . . .	11
1.4	Results of chloride calibration of Green Bay dispersive exchange . . . . .	13
1.5	Suspended particle calibration . . . . .	14
1.6	Organic carbon fraction of surface water particles . . . . .	15
1.7	Partitioning model: Sensitivity of dissolved chemical fraction to non-settling organic matter binding efficiency . . . . .	16
1.8	Calibration of partitioning model: Comparison to distribution coefficient data for HOCs in the Great Lakes . . . . .	17
1.9	Sensitivity of computed volatilization rate to wind speed and temperature (pentachlorobiphenyl) . . . . .	18
1.10	Plutonium deposition to Lake Michigan . . . . .	20
1.11	Total loading of lead to Lake Michigan . . . . .	20
1.12	PCBs loading time function and comparison to reported PCBs load estimates . . . . .	22
1.13	MICHTOX food chain structure . . . . .	24
1.14	Chemical assimilation efficiency for <i>Mysis</i> calculated from Lake Ontario PCBs data . . . . .	26
1.15	Chemical assimilation efficiency for alewife calculated from Lake Ontario HOCs data . . . . .	26
1.16	Chemical assimilation efficiency for <i>Diporeia</i> calculated from Lake Ontario PCBs data . . . . .	26
1.17	Steady-state spreadsheet model for pentachlorobiphenyl (PCB5) . . . . .	27
1.18	Simplified Lake Michigan lake trout food chain . . . . .	28



---

1.19	MICHTOX simulation of plutonium in southern Lake Michigan (epilimnion, hypolimnion, overturn data) . . . . .	31
1.20	MICHTOX simulation of plutonium in southern Lake Michigan (epilimnion, hypolimnion, and seasonal epilimnetic data) . . . . .	31
1.21	MICHTOX simulation of plutonium in southern Lake Michigan – sensitivity to vertical segmentation . . . . .	32
1.22	Simulation of annual-averaged lead concentrations in southern Lake Michigan . . . . .	32
1.23	Lead simulation in southern Lake Michigan surficial sediment . . . . .	32
1.24	Simulation of PCBs in southern Lake Michigan . . . . .	33
1.25	Simulation of PCBs in Green Bay . . . . .	33
1.26	PCBs simulations in central Lake Michigan and outer Green Bay . . . . .	34
1.27	PCBs simulations in Lake Michigan and outer Green Bay surficial sediments . . . . .	34
1.28	PCBs simulations in Green Bay sediments . . . . .	34
1.29	Simulated PCBs distribution in southern Lake Michigan sediments . . . . .	35
1.30	PCBs concentrations in Lake Michigan lake trout . . . . .	35
1.31	Verification of PCBs accumulation in age seven lake trout . . . . .	36
1.32	Sensitivity of trout PCBs predictions to the food chain . . . . .	36
1.33	Verification of PCBs accumulation predictions in lake trout age classes 2-9 . . . . .	37
1.34	Simulation of PCBs concentrations in age 2-12 trout, 1980-1989 . . . . .	37
1.35	Simulation of PCBs concentrations in age 2-12 trout, 1980-1990 . . . . .	37
1.36	Verification of PCBs concentration predictions for bloater . . . . .	38
1.37	Validation of bioaccumulation predictions for lower trophic levels in Lake Michigan . . . . .	39
1.38	MICHTOX predicted trout bioaccumulation and data from Lake Ontario . . . . .	40
1.39	MICHTOX predicted trout biota-to-sediment ratio and data from Lake Ontario . . . . .	41
1.40	Load-concentration relationship for PCBs in southern Lake Michigan . . . . .	42
1.41	Relative magnitude of PCBs fate and transport fluxes . . . . .	43
1.42	PCBs air-water fluxes at steady-state in southern Lake Michigan . . . . .	44

---

1.43	Sensitivity of water concentrations to $K_{ow}$ . . . . .	44
1.44	Sensitivity of trout concentrations to $K_{ow}$ . . . . .	44
1.45	Sensitivity to Henry's constant . . . . .	45
1.46	Sensitivity to volatilization rate . . . . .	45
1.47	Sensitivity to dry deposition velocity . . . . .	45
1.48	Sensitivity to sediment-water diffusion coefficient . . . . .	45
1.49	Sensitivity of trout concentration to chemical assimilation efficiency . . . . .	46
1.50	Sensitivity of trout concentration to pelagic diet fraction of forage fish . . . . .	46
1.51	Sensitivity to dispersive exchange coefficient . . . . .	46
1.52	Sensitivity to suspended particle concentration . . . . .	46
1.53	Sensitivity to particle burial velocity . . . . .	47
1.54	Sensitivity to particle settling velocity . . . . .	47
1.55	Sensitivity of water concentrations to suspended particle $f_{oc}$ . . . . .	47
1.56	Sensitivity of sediment concentrations to suspended particle $f_{oc}$ . . . . .	47
1.57	Sensitivity of trout concentrations to suspended particle $f_{oc}$ . . . . .	48
1.58	MICHTOX steady-state response to different tributary load distributions . . . . .	48
1.59	Load-concentration relationship for PCBs in southern Lake Michigan including confidence limits . . . . .	53
1.60	Load-concentration relationship for PCBs in southern Lake Michigan water: Effect of constant air concentration . . . . .	55
1.61	Load-concentration relationship for PCBs in southern Lake Michigan trout: Effect of constant air concentration . . . . .	55
1.62	Contribution of critical parameters to steady-state model uncertainty: Water concentration . . . . .	55
1.63	Contribution of critical parameters to steady-state model uncertainty: Trout concentration . . . . .	56
1.64	Predicted PCBs time response to loading reduction in southern Lake Michigan . . . . .	58
1.65	Response of trout PCBs concentrations at various times after reducing PCBs load . . . . .	58

---

1.66	Simulation of chlordane in Lake Michigan trout (Michigan Department of Natural Resources, 1990) . . . . .	59
1.67	Simulation of DDT in Lake Michigan trout (DeVault <i>et al.</i> , 1986; Michigan Department of Natural Resources, 1990) . . . . .	59
1.68	Simulation of dieldrin in Lake Michigan trout (DeVault <i>et al.</i> , 1986; Michigan Department of Natural Resources, 1990) . . . . .	59
1.69	Load cutoff simulation of PCBs in Lake Michigan trout . . . . .	59
1.70	Tetrachlorodibenzo-p-dioxin simulation in Lake Michigan trout (U.S. Environmental Protection Agency, 1989b) . . . . .	60
1.71	Tetrachlorodibenzofuran simulation in Lake Michigan trout (U.S. Environmental Protection Agency, 1989b) . . . . .	60
1.72	Predicted effectiveness of PCBs load reductions . . . . .	61
1.73	Simulation of storm event in southern Lake Michigan . . . . .	62
1.74	Effect of storm event on PCBs in southern Lake Michigan sediment . . . . .	62
1.75	Sensitivity of PCBs concentrations in trout to thin (1.1 cm) surficial sediment layer thickness . . . . .	63
1.76	Predicted water PCBs concentrations for ten realizations of dynamic model . . . . .	63
1.77	Predicted trout PCBs concentrations for ten realizations of dynamic model . . . . .	64
2.1	MICHTOX mass balance schematic . . . . .	87
2.2	Spatial segmentation for the 17 segment MICHTOX model . . . . .	88
2.3	The LMMBP estimates of total PCBs atmospheric vapor concentrations processed as monthly values for MICHTOX Segments 1-3 . . . . .	91
2.4	The LMMBP estimates of total PCBs atmospheric wet and dry deposition processed as monthly values for MICHTOX Segment 1 (southern Lake Michigan) . . . . .	92
2.5	The LMMBP estimates of total PCBs tributary loading processed as monthly values for MICHTOX Segment 1 (southern Lake Michigan) . . . . .	93
2.6	Long-term estimates of Lake Michigan total PCBs vapor concentrations . . . . .	99
2.7	Long-term estimates of Lake Michigan total PCBs atmospheric deposition loading . . . . .	100
2.8	Long-term estimates of Lake Michigan total PCBs tributary loading . . . . .	101

---

2.9	Comparison of MICHTOX steady-state total PCBs concentrations to Saugatuck fish data . . . . .	102
2.10	Comparison of MICHTOX steady-state total PCBs concentrations to Sheboygan Reef fish data . . . . .	103
2.11	Original MICHTOX predictions and data for main lake total PCBs concentrations . . . . .	105
2.12	Original MICHTOX predictions and data for main lake sediment total PCBs concentrations (sediment cores collected in 1991-1992) . . . . .	106
2.13	Original MICHTOX predictions of total PCBs concentrations in fish and comparison to Sheboygan Reef zone data . . . . .	107
2.14	Original MICHTOX predictions of total PCBs concentrations in lake trout and comparison to DeVault <i>et al.</i> (1986) data. . . . .	108
2.15	Long-term Scenario A predictions of main lake total PCBs concentrations . . . . .	111
2.16	MICHTOX southern Lake Michigan total PCBs predictions. Comparison of long-term Scenario A to original model predictions. . . . .	112
2.17	Comparison of long-term Scenario A predictions to main lake sediment total PCBs concentrations (sediment cores collected in 1991-1992) . . . . .	113
2.18	Comparison of long-term Scenario A predictions to DeVault <i>et al.</i> (1986) lake trout data. . . . .	114
2.19	Comparison of MICHTOX Scenario A total PCBs concentrations to Sheboygan Reef data . . . .	115
2.20	Long-term Scenario B predictions of main lake total PCBs concentrations . . . . .	117
2.21	MICHTOX southern Lake Michigan total PCBs predictions. Comparison of long-term Scenario B to original model predictions . . . . .	118
2.22	Comparison of long-term Scenario B predictions to the LMMBP deepwater dissolved total PCBs concentrations . . . . .	119
2.23	Comparison of Scenario B predictions to main lake sediment total PCBs concentrations (sediment cores collected in 1991-1992) . . . . .	120
2.24	Comparison of long-term Scenario B predictions to average total PCBs sediment concentrations (LMMBP and GBMBP box core samples) . . . . .	121
2.25	Comparison of long-term Scenario B predictions to DeVault <i>et al.</i> (1986) lake trout data . . . . .	122
2.26	Comparison of long-term Scenario B total PCBs concentrations to Sheboygan Reef fish data . . . . .	123
2.27	Long-term Scenario C predictions of main lake total PCBs concentrations. . . . .	124

---

2.28	MICHTOX southern Lake Michigan total PCBs predictions. Comparison of long-term Scenario C to original model predictions. ....	125
2.29	Comparison of Scenario C predictions to main lake sediment total PCBs concentrations (sediment cores collected in 1991-1992) ....	126
2.30	Comparison of long-term Scenario C predictions to DeVault <i>et al.</i> (1986) lake trout data ....	127
2.31	Comparison of MICHTOX Scenario C total PCBs concentrations to Sheboygan Reef fish data ....	128
2.32	Toxic chemical management alternatives. Comparison of forecast simulation for age seven lake trout in southern Lake Michigan ....	130
2.33	Total PCBs concentrations in Lake Michigan air and water ....	133
2.34	Ratio of total PCBs concentrations between dissolved water and vapor (MICHTOX No-Action forecast) ....	134
2.35	Sensitivity of food chain model to initial conditions. Age seven lake trout: No-Change forecast ....	135
2.36	Sensitivity of MICHTOX steady-state PCBs concentrations to chemical assimilation efficiency and comparison to Saugatuck biota zone data ....	137
2.37	Predicted bioaccumulation factors for total PCBs in age seven lake trout (No-Change forecast simulation) ....	138

## List of Tables

1.1	Sediment Segment Parameterization. . . . .	13
1.2	Particle Flux Parameterization . . . . .	14
1.3	Selected Air Concentrations and Calculated Atmospheric Deposition Loadings for Lake Michigan Priority Pollutants . . . . .	19
1.4	Octanol-Water Partition Coefficient ( $K_{ow}$ ) and Organic Carbon Partition Coefficient ( $K_{oc}$ ) for Lake Michigan Priority Toxics . . . . .	29
1.5	Physicochemical Parameters Used in Volatilization Parameterization of Lake Michigan Critical Pollutants . . . . .	29
1.6	Volatilization Rate Parameters for Lake Michigan Critical Pollutants . . . . .	30
1.7	Photolysis Rate for Lake Michigan Critical Pollutants . . . . .	30
1.8	Predicted Steady-State Bioaccumulation Factors (BAFs) for Critical Pollutants in Lake Michigan Lake Trout . . . . .	40
1.9	Predicted Steady-State Concentrations of Critical Pollutants in Lake Michigan for Unit Loading. . . . .	42
1.10	Summary of Mass Fate and Transport for Critical Pollutants in Lake Michigan. Southern Lake Michigan Mass Fluxes Expressed as Fractions of Segment Load . . . . .	43
1.11	Comparison of PCBs Mass Fate and Transport for Critical Pollutants in Southern Lake Michigan and Central Green Bay . . . . .	43
1.12	Probability Distribution for Steady-State Model Uncertainty Analysis . . . . .	51
1.13	Summary of Results for First Test of Model Uncertainty. Predicted Steady-State Concentrations for Fixed Air Concentrations of 1 ng/m <sup>3</sup> for Each Chemical. . . . .	52
1.14	Summary of Results for Second Test of Model Uncertainty. Predicted Steady-State Concentrations for Expected Air Concentrations of Each Chemical. . . . .	53

---

1.15	Summary of Results for Third Test of Model Uncertainty. Predicted Steady-State PCBs Concentrations for Varying Tributary Loads and Expected Air Concentrations . . . . .	54
1.16	Predicted Long-Term Chemical Loss Rates for Lake Michigan Critical Pollutants Following Loading Reduction. First-Order Loss Rates in Units of 1/Year in Southern Lake Michigan . . . . .	57
2.1	Cruise- and Segment-Specific Dissolved Fraction of Total PCBs Concentrations (ng/L) . . . . .	94
2.2	Cruise- and Segment-Specific Average Particulate Total PCBs Concentrations (ng/L) . . . . .	94
2.3	Segment-Specific Average Surficial Sediment Total PCBs Concentrations (ng/g) . . . . .	95
2.4a	Average Total PCBs Concentrations in Fish in the Saugatuck Biota Zone. . . . .	95
2.4b	Average Total PCBs Concentrations in Fish in the Sheboygan Reef Biota Zone . . . . .	96
2.4c	Average Total PCBs Concentrations in Fish in the Sturgeon Bay Biota Zone . . . . .	96
2.5	Comparison of Original MICHTOX Total PCBs Forcing Functions for 1994-1995 to the LMMBP Estimates (Whole-Lake Average) . . . . .	104
2.6	Mass Balance Diagnostics for Total PCBs in the Original MICHTOX Simulation (Year 1994-1995) . . . . .	109
2.7	Comparison of Original and Revised Henry's Constant ( $\text{atm m}^3/\text{mol}$ ) Parameterization . . . . .	110
2.8	Monthly PCBs Volatilization Rates (m/d) Calculated by Original and Revised MICHTOX Formulations . . . . .	111
2.9	Mass Balance Diagnostics for Total PCBs in MICHTOX Scenario B Simulation (Year 1994-1995) . . . . .	127
2.10	MICHTOX Predictions of Total PCBs Concentrations ( $\mu\text{g/g}$ ) in Lake Trout for Toxic Chemical Management Alternatives . . . . .	131
2.11	Results of BMC Uncertainty Analysis for Original Steady-State MICHTOX Model . . . . .	132

---

## Abbreviations

AOCs	Areas of Concern
BAF	Bioaccumulation factor
BaP	Benzo(a)pyrene
BCF	Bioconcentration factor
BMC	Bayesian Monte Carlo
BSF	Biota-to-sediment factor
BSR	Biota-to-sediment ratio
CV	Coefficient of variance
DDD	Dichlorodiphenyldichloroethane
DDE	Dichlorodiphenyldichloroethylene
DDT	Dichlorodiphenyltrichloroethane
EPRI	Electric Power Research Institute
ERL	Environmental Research Laboratory
GBMBP	Green Bay Mass Balance Project
GLEC	Great Lakes Environmental Center
GLNPO	Great Lakes National Program Office
HCB	Hexachlorobenzene
HOCs	Hydrophobic organic chemicals
LaMP	Lake-wide Management Plan
LCL	Lower confidence limit
LMMBP	Lake Michigan Mass Balance Project
LLRFRB	Large Lakes and Rivers Forecasting Research Branch
LLRS	Large Lakes Research Station
lnCV	Lognormal coefficient of variance
MDNR	Michigan Department of Natural Resources
MED	Mid-Continent Ecology Division
NSOM	Non-settling organic matter
PCBs	Polychlorinated biphenyls
PCB4	Tetrachlorobiphenyl
PCB5	Pentachlorobiphenyl
PCDD	Pentachlorodibenzo-p-dioxin
PCDF	Pentachlorodibenzofuran
QAPP	Quality Assurance Project Plan
TCDD	Tetrachlorodibenzo-p-dioxin
TCDF	Tetrachlorodibenzofuran
UCL	Upper confidence limit
USEPA	United States Environmental Protection Agency
USFWS	United States Fish and Wildlife Service
VWA	Volume-weighted average



---

## **Acknowledgments**

Special thanks for the assistance and cooperation of the federal and on-site contractor staff at the U.S. Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Mid-Continent Ecology Division-Duluth, Large Lakes Research Station, Grosse Ile, Michigan. David Griesmer, Xiangsheng Xia, Katie Taunt, Xiaomi Zhang, and Xin Zhang provided raw and processed Lake Michigan Mass Balance Project and historical data used for the modeling. Wilson Melendez provided programming support for the necessary modifications of MICHTOX mass balance, food web bioaccumulation, and post-processing programs. Thanks to Kay Morrison for the graphic renditions and figures and to Debra Caudill for formatting and word processing.

---

## Executive Summary

MICHTOX is a toxic chemical mass balance and food chain bioaccumulation model that was first developed in the early 1990s. A Bayesian Monte Carlo uncertainty analysis demonstrated that MICHTOX predicted polychlorinated biphenyl (PCBs) concentrations should be within a factor of two of the measured data. During the early part of the Lake Michigan Mass Balance Project (LMMBP), MICHTOX was updated and used as a preliminary assessment tool of the LMMBP PCBs data and to provide a screening-level analysis of the potential future trends in total PCBs concentrations in Lake Michigan water, sediment, and fish under a variety of contaminant load scenarios.

As reported in 1992, the model predicted the response of Lake Michigan, and with additional resolution, Green Bay to atmospheric and tributary loadings. With its bioaccumulation component, chemical accumulation in biota was predicted in response to the loadings. The model is capable of either dynamic or steady-state simulations. Dynamic model predictions were used to predict the long-term rate of concentration decline following load reduction for each toxic chemical. Significant reductions of PCBs in lake trout were predicted for 2000 with no additional loading reductions. Additional reductions of PCBs concentrations could only be achieved with significant reductions in atmospheric sources. These results were uncertain because PCBs loading history is poorly defined and because of potential error in the parameterization of the surficial sediment layer thickness. The thickness of this layer was demonstrated to be a critical factor in model uncertainty. Additional factors leading to model uncertainty included uncertainty in initial concentrations and loading history and dynamics of the Lake Michigan trophic structure.

As reported in 2002, MICHTOX was used to provide a preliminary mass balance modeling assessment of PCBs in Lake Michigan. Because PCBs vapor concentrations from the LMMBP were significantly higher than estimated in the original model, total PCBs forcing functions were recalculated using the LMMBP estimates. Recommended changes to the model increased the volatilization mass transport rates, resulting in the PCBs equilibrium shifting significantly towards the atmospheric vapor phase quicker than previously predicted. This demonstrated that air-water fluxes predominated the transport pathways for PCBs in Lake Michigan. The best prediction of PCBs concentrations in water, sediment, and fish were obtained with the forcing function peaking in 1961-1963. This was different than the original model simulation reported in 1992. The model was used to forecast total PCBs concentrations in lake trout for a variety of scenarios representing alternative strategies for managing PCBs in Lake Michigan. Because of model uncertainty, observed average total PCBs concentrations should be within a factor of two of predicted values. The bioaccumulation predictions were not sensitive to initial conditions but were sensitive to model parameterization. The PCB predictions of this model are historic and have been replaced by the predictions derived from the improved models used for the LMMBP.